

# Unit distances between disjoint convex translates

27 April 2026

## Abstract

Let  $h(n)$  be the maximum number of pairs at Euclidean set-distance exactly 1 among  $n$  pairwise disjoint translates of one compact convex set in the plane. Erdős and Pach proved the upper bound  $h(n) = O(n^{4/3})$ . We give a self-contained matching lower-bound construction. Combining the two bounds gives the asymptotic order

$$h(n) = \Theta(n^{4/3}).$$

In particular, Erdős Problem #956 has a positive answer in the requested polynomial-growth sense: one may take any constant  $c < 1/3$  in  $h(n) > n^{1+c}$  for all sufficiently large  $n$ .

## 1 Reduction to a difference body

For compact sets  $A, B \subset \mathbb{R}^2$ , write

$$\delta(A, B) = \inf\{\|a - b\|_2 : a \in A, b \in B\}.$$

Let  $C \subset \mathbb{R}^2$  be compact and convex, and put

$$D := C - C = \{c - c' : c, c' \in C\}.$$

Then  $D$  is centrally symmetric and convex. For two translation vectors  $x, y \in \mathbb{R}^2$ ,

$$\delta(C + x, C + y) = \text{dist}(y - x, D), \tag{1}$$

where the distance on the right is ordinary Euclidean point-to-set distance. Moreover,

$$(C + x) \cap (C + y) = \emptyset \iff y - x \notin D. \tag{2}$$

Conversely, every centrally symmetric compact convex set  $D$  is of the form  $C - C$ , namely with  $C = D/2$ : by central symmetry,  $-D = D$ , and by convexity  $D + D = 2D$ , so

$$(D/2) - (D/2) = \frac{1}{2}(D + (-D)) = \frac{1}{2}(D + D) = D.$$

Thus it is enough to construct a centrally symmetric compact convex body  $D$  and a finite point set  $X$  such that no non-zero difference from  $X - X$  lies in  $D$ , while many differences have Euclidean distance exactly 1 from  $D$ .

## 2 A small convex body whose unit offset contains a parabolic arc

Fix parameters

$$0 < W \leq 1, \quad \eta := W^4.$$

For  $0 \leq t \leq W$ , define

$$\gamma(t) := \left( t, 1 + \eta - \frac{t^2}{2} \right), \quad \nu(t) := \frac{(t, 1)}{\sqrt{1+t^2}},$$

and

$$p(t) := \gamma(t) - \nu(t).$$

Thus  $\nu(t)$  is the upward unit normal to the parabola  $\gamma$ , and  $p(t)$  is the point obtained by moving one unit inward from  $\gamma(t)$  along this normal. Let

$$D_W := \text{conv}(\{\pm p(t) : 0 \leq t \leq W\}). \quad (3)$$

This is a centrally symmetric compact convex body for every  $W > 0$ : it is compact and convex by definition, centrally symmetric by construction, and the three points  $p(0)$ ,  $-p(0)$ , and  $p(W)$  are not collinear.

**Lemma 1.** For  $0 \leq t \leq W \leq 1$ ,

$$0 \leq p_x(t) \leq \frac{W^3}{2}, \quad 0 \leq p_y(t) \leq \eta.$$

Consequently

$$D_W \subseteq \left[ -\frac{W^3}{2}, \frac{W^3}{2} \right] \times [-\eta, \eta]. \quad (4)$$

*Proof.* Write  $r(t) = \sqrt{1+t^2}$ . Then

$$p_x(t) = t \left( 1 - \frac{1}{r(t)} \right) \geq 0,$$

and, since

$$1 - \frac{1}{\sqrt{1+u}} = \frac{u}{\sqrt{1+u}(\sqrt{1+u}+1)} \leq \frac{u}{2} \quad (u \geq 0),$$

we have  $p_x(t) \leq t^3/2 \leq W^3/2$ .

For the  $y$ -coordinate,

$$p_y(t) = \eta + 1 - \frac{t^2}{2} - \frac{1}{\sqrt{1+t^2}}.$$

We use the elementary estimates

$$1 - \frac{u}{2} \leq \frac{1}{\sqrt{1+u}} \leq 1 - \frac{u}{2} + \frac{u^2}{2} \quad (0 \leq u \leq 1). \quad (5)$$

The left inequality follows, for instance, from  $\sqrt{1+u} \leq 1 + u/2$  and  $1/(1+u/2) \geq 1 - u/2$ . For the right inequality, let

$$F(u) := 1 - \frac{u}{2} + \frac{u^2}{2} - (1+u)^{-1/2}.$$

Then  $F(0) = F'(0) = 0$  and

$$F''(u) = 1 - \frac{3}{4}(1+u)^{-5/2} \geq \frac{1}{4} \quad (u \geq 0),$$

so  $F(u) \geq 0$ . Applying (5) with  $u = t^2$  gives

$$p_y(t) \leq \eta$$

and

$$p_y(t) \geq \eta - \frac{t^4}{2} \geq W^4 - \frac{W^4}{2} \geq 0.$$

The box containment follows from the definition of  $D_W$  as the convex hull of these points and their negatives.  $\square$

**Lemma 2.** *For every  $0 \leq t \leq W$ ,*

$$\text{dist}(\gamma(t), D_W) = 1.$$

*Proof.* Since  $p(t) \in D_W$  and  $\gamma(t) - p(t) = \nu(t)$  is a unit vector, we have

$$\text{dist}(\gamma(t), D_W) \leq 1.$$

It remains to prove the reverse inequality.

We show that the line

$$\{z \in \mathbb{R}^2 : \langle z, \nu(t) \rangle = \langle p(t), \nu(t) \rangle\}$$

supports  $D_W$ . First consider another point  $p(s)$ ,  $0 \leq s \leq W$ . The vertical shift by  $\eta$  cancels in the following calculation, so put  $r_u := \sqrt{1+u^2}$ . A direct computation gives

$$\begin{aligned} & \langle p(t), \nu(t) \rangle - \langle p(s), \nu(t) \rangle \\ &= \frac{(s-t)^2}{2r_t} - \left(1 - \frac{1+st}{r_s r_t}\right). \end{aligned}$$

Furthermore,

$$1 - \frac{1+st}{r_s r_t} = \frac{(s-t)^2}{r_s r_t (r_s r_t + 1 + st)}.$$

Since  $s, t \geq 0$ , we have

$$r_s (r_s r_t + 1 + st) \geq 2.$$

Therefore

$$\frac{(s-t)^2}{2r_t} \geq \frac{(s-t)^2}{r_s r_t (r_s r_t + 1 + st)},$$

and hence

$$\langle p(s), \nu(t) \rangle \leq \langle p(t), \nu(t) \rangle. \quad (6)$$

By Lemma 1, both coordinates of every  $p(s)$  are non-negative, and both coordinates of  $\nu(t)$  are non-negative. Hence

$$\langle -p(s), \nu(t) \rangle \leq 0 \leq \langle p(t), \nu(t) \rangle.$$

Together with (6), this proves that every generator  $\pm p(s)$  of  $D_W$  lies in the closed half-plane

$$\langle z, \nu(t) \rangle \leq \langle p(t), \nu(t) \rangle.$$

Thus the same is true for all  $z \in D_W$ . Consequently,

$$\text{dist}(\gamma(t), D_W) \geq \langle \gamma(t), \nu(t) \rangle - \langle p(t), \nu(t) \rangle = \langle \gamma(t) - p(t), \nu(t) \rangle = 1.$$

This proves the lemma.  $\square$

### 3 The point set

Fix a small absolute constant  $\alpha \in (0, 1/10]$ . For an integer  $k \geq 2$ , put

$$W := \frac{\alpha}{k}, \quad a := \frac{W}{k} = \frac{\alpha}{k^2}, \quad b := \frac{a^2}{2} = \frac{\alpha^2}{2k^4}, \quad \eta := W^4 = \frac{\alpha^4}{k^4}.$$

Let  $D = D_W$  be the body from (3). Define two grids

$$L := \{(ra, sb) : 0 \leq r \leq k, 0 \leq s \leq k^2\},$$

$$U := \{(ra, 1 + \eta + sb) : 0 \leq r \leq k, 0 \leq s \leq k^2\},$$

where  $r, s$  are integers. Set

$$X_k := L \cup U.$$

Then

$$n_k := |X_k| = 2(k+1)(k^2+1) = 2k^3 + O(k^2). \quad (7)$$

**Lemma 3.** *No non-zero difference of two points of  $X_k$  lies in  $D$ .*

*Proof.* By Lemma 1,

$$D \subseteq [-W^3/2, W^3/2] \times [-\eta, \eta].$$

Since  $W = \alpha/k$  and  $a = \alpha/k^2$ ,

$$\frac{W^3}{2} = \frac{\alpha^3}{2k^3} < \frac{\alpha}{k^2} = a$$

for all  $k \geq 2$  and  $\alpha \leq 1$ . Also,

$$\eta = \frac{\alpha^4}{k^4} < \frac{\alpha^2}{2k^4} = b$$

because  $\alpha^2 < 1/2$ .

If two points lie in the same grid  $L$  or  $U$ , then their difference is of the form  $(ma, \ell b)$  with integers  $m, \ell$ . If this difference is non-zero, then either  $|ma| \geq a > W^3/2$  or  $|\ell b| \geq b > \eta$ , so it is not in  $D$ .

If one point lies in  $L$  and the other in  $U$ , then the absolute value of the  $y$ -coordinate of their difference is at least

$$1 + \eta - k^2 b = 1 + \eta - \frac{W^2}{2} \geq 1 - \frac{\alpha^2}{2} > \eta,$$

again excluding membership in  $D$ . Thus no non-zero difference from  $X_k - X_k$  lies in  $D$ .  $\square$

**Lemma 4.** *There are at least*

$$M_k := \sum_{i=1}^k (k+1-i)(k^2+1-i^2) = \frac{5}{12}k^4 + O(k^3)$$

*unordered pairs  $\{x, y\} \subset X_k$  such that*

$$\text{dist}(y - x, D) = 1.$$

*Proof.* For  $i = 1, 2, \dots, k$ , put

$$t_i := ia.$$

Then  $0 < t_i \leq W$ , and

$$\gamma(t_i) = \left( ia, 1 + \eta - \frac{i^2 a^2}{2} \right) = (ia, 1 + \eta - i^2 b).$$

By Lemma 2,  $\text{dist}(\gamma(t_i), D) = 1$ .

Now choose integers

$$0 \leq r \leq k - i, \quad i^2 \leq s \leq k^2.$$

Then

$$(ra, sb) \in L, \quad ((r + i)a, 1 + \eta + (s - i^2)b) \in U,$$

and their difference is exactly

$$(ia, 1 + \eta - i^2 b) = \gamma(t_i).$$

Thus each such choice gives a pair at distance 1 from  $D$ . These counted pairs are distinct, since each has one endpoint in  $L$ , one endpoint in  $U$ , and horizontal separation  $ia$ . For fixed  $i$ , the number of choices is

$$(k + 1 - i)(k^2 + 1 - i^2).$$

Summing over  $i = 1, \dots, k$  gives  $M_k$  pairs. The displayed closed form follows from the elementary sums

$$\sum_{i=1}^k i = \frac{k(k+1)}{2}, \quad \sum_{i=1}^k i^2 = \frac{k(k+1)(2k+1)}{6}, \quad \sum_{i=1}^k i^3 = \frac{k^2(k+1)^2}{4}.$$

Indeed,

$$M_k = \frac{5}{12}k^4 + \frac{1}{6}k^3 + \frac{1}{12}k^2 + \frac{1}{3}k.$$

□

## 4 The lower bound for convex translates

**Theorem 1.** *There is an absolute constant  $c_0 > 0$  such that*

$$h(n) \geq c_0 n^{4/3}$$

for all sufficiently large  $n$ .

*Proof.* For the point set  $X_k$  and body  $D = D_W$  above, let

$$C := \frac{1}{2}D.$$

Since  $D$  is centrally symmetric and convex,  $C - C = D$ . Lemma 3 and (2) imply that the translates  $C + x$ ,  $x \in X_k$ , are pairwise disjoint. Lemma 4 and (1) imply that at least  $M_k$  pairs among them have set-distance exactly 1.

By (7),  $n_k = 2k^3 + O(k^2)$ , while Lemma 4 gives  $M_k = (5/12)k^4 + O(k^3)$ . Hence

$$h(n_k) \geq M_k \geq c_1 n_k^{4/3}$$

for an absolute constant  $c_1 > 0$  and all sufficiently large  $k$ .

For a general sufficiently large  $n$ , choose  $k$  maximal with  $n_k \leq n$ . Add  $n - n_k$  extra translates very far from all existing translates and from one another. More explicitly, since  $D$  and  $X_k$  are bounded, one may choose a vector  $v$  with  $\|v\|$  larger than  $\sup_{d \in D} \|d\| + \sup_{x \in X_k} \|x\|$  and add suitable points among  $v, 2v, 3v, \dots$ . Then every new difference has norm larger than  $\sup_{d \in D} \|d\|$ , and hence is outside  $D$ . This preserves disjointness and cannot decrease the number of distance-1 pairs. Since  $n_{k+1}/n_k \rightarrow 1$ , the same lower bound, with a smaller absolute constant, holds for all sufficiently large  $n$ .  $\square$

## 5 Asymptotic determination of $h(n)$

We use the following published theorem of Erdős and Pach.

**Theorem 2** (Erdős–Pach [2]). *For the same function  $h(n)$ ,*

$$h(n) = O(n^{4/3}).$$

This is also the upper bound recorded on Erdős Problem #956 [1]. Combining Theorem 2 with Theorem 1 gives the asymptotic order.

**Corollary 1.** *For the maximum number  $h(n)$  of unit distances between  $n$  pairwise disjoint translates of one compact convex plane set,*

$$h(n) = \Theta(n^{4/3}).$$

*Consequently, for every fixed  $c < 1/3$  there is  $n_0(c)$  such that*

$$h(n) > n^{1+c} \quad (n \geq n_0(c)).$$

*Remark 1* (Relation to Valtr’s parabolic norm construction). The local counting mechanism above is the same parabolic-grid mechanism used by Valtr to construct strictly convex norms with many unit distances [3]. The additional point in this note is that the convex set  $D$  is made extremely small: it is essentially the inner unit parallel body of a tiny parabolic cap. This makes the translates of  $C = D/2$  pairwise disjoint, while the outer unit parallel boundary still contains the required parabolic difference vectors.

## References

- [1] T. F. Bloom, *Erdős Problem #956*, <https://www.erdosproblems.com/956>, accessed 27 April 2026.
- [2] P. Erdős and J. Pach, *Variations on the theme of repeated distances*, *Combinatorica* **10** (1990), no. 3, 261–269. doi:10.1007/BF02122780.
- [3] P. Valtr, *Strictly convex norms allowing many unit distances and related touching questions*, manuscript, 2005. Available at <https://kam.mff.cuni.cz/~valtr/n.pdf>.